



Daffodil International University

Faculty of Science & Information Technology
Department of Computer Science & Engineering

Final Examination, Spring 2026

Course Code: CSE226, Course Title: Numerical Methods

Level: 2 Term: 3, Batch: 66

Time: 2:00 hrs.

Marks: 40

Answer All Questions

[The figures in the right margin indicate the full marks and corresponding course outcomes. All portions of each question must be answered sequentially]

question must be answered sequentially

1.	(a) Apply Gauss-Jacobi method to solve the following system of linear equations until converges correct to 3 decimal places by considering initial guess $x_0 = y_0 = z_0 = 0$. $\begin{aligned} 4x - 10y + 3z &= -3 \\ x + 6y + 10z &= -3 \\ 10x - 5y - 2z &= 3 \end{aligned}$	[7]	CO2											
	(b) Identify the first derivative of $x=1.1$ from the following table <table border="1" style="margin-left: auto; margin-right: auto; border-collapse: collapse;"> <tr> <td style="padding: 5px 10px;">x</td> <td style="padding: 5px 10px;">1.0</td> <td style="padding: 5px 10px;">1.1</td> <td style="padding: 5px 10px;">1.2</td> <td style="padding: 5px 10px;">1.3</td> <td style="padding: 5px 10px;">1.4</td> </tr> <tr> <td style="padding: 5px 10px;">y</td> <td style="padding: 5px 10px;">7.989</td> <td style="padding: 5px 10px;">8.403</td> <td style="padding: 5px 10px;">8.781</td> <td style="padding: 5px 10px;">9.129</td> <td style="padding: 5px 10px;">9.451</td> </tr> </table>	x		1.0	1.1	1.2	1.3	1.4	y	7.989	8.403	8.781	9.129	9.451
x	1.0	1.1	1.2	1.3	1.4									
y	7.989	8.403	8.781	9.129	9.451									
2.	Apply Runge-Kutta method of 4th order to calculate approximate value of $y(0.1)$ and $y(0.2)$ of $\frac{dy}{dx} = x - y^2$, $y(0)=1$ correct to 4 decimal places.	[10]												
3.	Identify the best method among Weddle's, Simpson's 1/3rd and Simpson's 3/8th to compare with the exact value, by solving the integral of $I = \int_{0.5}^{0.8} \sqrt{x} e^{-x} dx$	[10]												
4.	(a) Certain experimental data of x and y are given as (0,-1), (2, 5), (5, 12), (7, 20). If the straight line $y = a_0 + a_1x$ is fitted to the given data by applying least square method, find out the approximate values of a_0 and a_1.	[4]	CO3											
	(b) Develop the second degree polynomial by using least square method for the following data: <table border="1" style="margin-left: auto; margin-right: auto; border-collapse: collapse;"> <tr> <td style="padding: 5px 10px;">x</td> <td style="padding: 5px 10px;">1</td> <td style="padding: 5px 10px;">3</td> <td style="padding: 5px 10px;">4</td> <td style="padding: 5px 10px;">6</td> </tr> <tr> <td style="padding: 5px 10px;">y</td> <td style="padding: 5px 10px;">0.63</td> <td style="padding: 5px 10px;">2.05</td> <td style="padding: 5px 10px;">4.08</td> <td style="padding: 5px 10px;">10.78</td> </tr> </table>	x		1	3	4	6	y	0.63	2.05	4.08	10.78	[6]	
x	1	3	4	6										
y	0.63	2.05	4.08	10.78										